

D($^{18}\text{F}, p\alpha$) ^{15}N reaction applied to nova γ -ray emission

N. de Séréville,¹ A. Coc,¹ C. Angulo,² M. Assunção,¹ D. Beaumel,³ B. Bouzid,⁴ S. Cherubini,^{2,*}
M. Couder,² P. Demaret,² F. de Oliveira Santos,⁵ P. Figuera,⁶ S. Fortier,³ M. Gaelens,²
F. Hammache,^{7,†} J. Kiener,¹ A. Lefebvre,¹ D. Labar,² P. Leleux,² M. Loiselet,² A. Ninane,²
S. Ouichaoui,⁴ G. Ryckewaert,² N. Smirnova,⁸ V. Tatischeff,¹ and J.-P. Thibaud¹

¹CSNSM, IN2P3/CNRS and Université Paris-Sud, F-91405 Orsay Campus, France

²Université catholique de Louvain, Chemin du Cyclotron 2, B-1348 Louvain-la-Neuve, Belgium

³IPN, IN2P3/CNRS and Université Paris-Sud, F-91406 Orsay Cedex, France

⁴USTHB, B.P. 32, El-Alia, Bab Ezzouar, Algiers, Algeria

⁵GANIL, B.P. 5027, 14021 Caen Cedex, France

⁶Laboratori Nazionali del Sud, INFN, Via S. Sofia, 44 - 95123 Catania, Italy

⁷GSI mbH, Planckstr. 1, D-64291 Darmstadt, Germany

⁸Instituut voor Kern en Stralingsfysika, Celestijnenlaan 200D, B-3001, Leuven, Belgium

(Dated: February 8, 2008)

The $^{18}\text{F}(p,\alpha)^{15}\text{O}$ reaction is recognized to be one of the most important reactions for nova gamma-ray astronomy as it governs the early $E \leq 511$ keV gamma emission. However in the nova temperature regime, its rate remains largely uncertain due to unknown low-energy resonance strengths. We report here the measurement of the $\text{D}(^{18}\text{F}, p)^{19}\text{F}(\alpha)^{15}\text{N}$ one-nucleon transfer reaction, induced by a 14 MeV ^{18}F radioactive beam impinging on a CD_2 target; outgoing protons and ^{15}N (or α -particles) were detected in coincidence in two silicon strip detectors. A DWBA analysis of the data resulted in new limits to the contribution of low-energy resonances to the rate of the $^{18}\text{F}(p,\alpha)^{15}\text{O}$ reaction.

PACS numbers: 25.60.Je, 21.10.Jx, 26.30.+k, 27.20.+n

Gamma-ray emission from classical novae is dominated, during the first hours, by positron annihilation resulting from the beta decay of radioactive nuclei. The main contribution comes from the decay of ^{18}F (half-life of 110 min) and hence is directly related to ^{18}F formation during the outburst[1, 2, 3]. A good knowledge of the nuclear reaction rates of production and destruction of ^{18}F is required to calculate the amount of ^{18}F synthesized in novae and the resulting gamma-ray emission. The rate relevant for the main mode of ^{18}F destruction (i.e, through $^{18}\text{F}(p,\alpha)^{15}\text{O}$) has been the object of many experiments[4, 5] (see also refs. in [3]). However, this rate remains poorly known at novae temperatures (lower than 3.5×10^8 K) due to the scarcity of spectroscopic information for levels near the proton threshold in the compound nucleus ^{19}Ne . This uncertainty is directly related to the unknown proton widths (Γ_p) of the first three levels (E_x , $J^\pi = 6.419$ MeV, $3/2^+$; 6.437 MeV, $1/2^-$ and 6.449 MeV, $3/2^+$). The tails of the corresponding resonances (at respectively $E_R = 8, 26$ and 38 keV) can dominate the astrophysical S-factor in the relevant energy range [3]. As a consequence of these nuclear uncertainties, the ^{18}F production in novae and the early gamma-ray emission used to be uncertain by a factor of ≈ 300 [3]. Unfortunately, a direct measurement of the relevant resonance strengths is impossible due to the very low Coulomb bar-

rier penetrability. Hence, we used an indirect method aiming at the determination of the one nucleon spectroscopic factors (S) in the analog levels of the mirror nucleus (^{19}F) by a neutron transfer reaction: $\text{D}(^{18}\text{F}, p)^{19}\text{F}$. Assuming the equality of one nucleon spectroscopic factors in analog levels, it is possible to calculate the proton widths through the relation $\Gamma_p = S \times \Gamma_{s.p.}$ where $\Gamma_{s.p.}$ is the single particle width. In this rapid communication we present the experimental technique, the results of a DWBA analysis and the implications on the $^{18}\text{F}(p,\alpha)^{15}\text{O}$ thermonuclear reaction rate.

The experiment has been carried out at the *Centre de Recherches du Cyclotron* at Louvain-la-Neuve (Belgium) where we used a ^{18}F radioactive beam. The ^{18}F was produced through the $^{18}\text{O}(p,n)^{18}\text{F}$ reaction, chemically extracted to form CH_3^{18}F molecules, transferred to the cyclotron source, ionized to the 2^+ state and accelerated to 14 MeV[6]. A total of 15 bunches of $\lesssim 1$ Ci of ^{18}F were produced providing each a mean beam intensity of 2.2×10^6 particles per second over a period of ≈ 2 hours. The beam contamination from its stable isobar was very small, $^{18}\text{O} / ^{18}\text{F} \leq 10^{-3}$. The ^{18}F beam was directed on deuterated polyethylene (CD_2) targets of ≈ 100 $\mu\text{g}/\text{cm}^2$ thickness made at Louvain-La-Neuve. For the energy considered here (1.4 MeV in the center of mass), the deuteron and the outgoing proton are both below the Coulomb barrier. In a similar transfer reaction $^{19}\text{F}(d,p)^{20}\text{F}$ [7] studied at the same c.m. energy, strong direct contributions have been observed. Hence, with a stripping integral dominated by the best known portion of the wave-function, accurate extraction of spectroscopic factors is expected.

The experimental setup is depicted in Figure 1. It

*Present address: EP3, Ruhr-Universität-Bochum, Bochum, Germany.

†Permanent address: IPN, IN2P3/CNRS and Université Paris-Sud, F-91406 Orsay Cedex, France.

consists of two silicon multistrip detectors composed of sectors with 16 concentric strips[8]. The first detector, LAMP, is positioned 9 cm upstream from the target; it consists of 6 sectors forming a conical shape covering laboratory angles between 115° and 160° (i.e. forward center-of-mass angles between 12° and 40°) providing a good acceptance for protons in the domain of interest for the differential cross section. The other detector, LEDA, is made up of 8 sectors forming a disk positioned 40 cm downstream from the target and is used for background reduction and normalization. The levels of interests are situated high above the alpha emission threshold (at 4.013 MeV) and their almost exclusive decay mode is through $^{19}\text{F}^* \rightarrow ^{15}\text{N} + \alpha$. Hence, to reduce background, we required coincidences between a proton in LAMP and a ^{15}N (or an α -particle discriminated by time of flight) in LEDA. Following Monte-Carlo simulations, the proton detection efficiency is found to be 27% and is only slightly reduced to 19% when the coincidence condition is applied.




FIG. 1: Experimental setup.

Thanks to the kinematics, only protons and α from $\text{D}(^{18}\text{F}, \text{p})^{19}\text{F}$ and $\text{D}(^{18}\text{F}, \alpha)^{16}\text{O}$ can reach LAMP. Using energy, angle and time-of-flight information, the events of interest with one proton in LAMP are selected. The excitation energy of the decaying ^{19}F levels can be kinematically reconstructed from the energies and angles of the detected protons and the known beam energy. The energy calibration of the silicon strips has been done with a triple-line alpha source and a small correction ($\approx 1\%$) on the excitation energy was introduced using a linear fit to the well-known energies of some lines in $\text{D}(^{18}\text{F}, \text{p})^{19}\text{F}$. This correction is needed because the reconstructed energy is very sensitive to the precise detector position (a 1 mm error induces a ≈ 10 keV energy shift). The statistical error from the fitting procedure induced by this correction is about 5 keV at 6.5 MeV. The corresponding coincidence spectrum is shown in Figure 2 where vertical lines correspond to the known position of the ^{19}F

levels[9] populated with low transferred angular momentum ($l \leq 2$). The resolution (≈ 100 keV FWHM) is not sufficient to separate the various levels, but the two $3/2^+$ levels of interest at 6.497 and 6.528 MeV (the analogs of the $3/2^+$ levels in ^{19}Ne) are well separated from the other groups of levels.

The angular distribution shown in Figure 3 is the result of a selection made on the 6.5 MeV peak in the coincidence spectrum. The coincidence efficiency is determined with a Monte-Carlo simulation taking an isotropic angular distribution for the α -decay of the ^{19}F . Coincidences were initially designed to eliminate $^{18}\text{O}(\text{d}, \text{p})^{19}\text{O}$ events induced by a possible isobaric beam contamination. This was not essential for this purpose because of the high beam purity, but it was useful to reduce the background from electronic noise affecting the low proton energy region. We checked that the ratio between coincidence and single events was well reproduced by Monte-Carlo simulations. Rutherford elastic scattering of ^{18}F on carbon from the target, detected in LEDA, provides the (target thickness) $\times$ (beam intensity) normalization. At such low beam intensity, the target stoichiometry is not expected to be modified during the experiment. This was confirmed by proton elastic scattering analysis of the targets done after the experiment at the Orsay ARAMIS facility. The solid lines in Figure 3 correspond to theoretical DWBA calculations made with the code FRESKO[10] in the zero-range approximation for different transferred angular momentum ($l = 0, 2$). The nuclear potentials were taken from Ref. [7] in which a similar neutron transfer reaction, $^{19}\text{F}(\text{d}, \text{p})^{20}\text{F}$, was studied at the same center-of-mass energy (subcoulomb transfer). A typical direct mechanism pattern was observed for 11 angular distributions of ^{20}F excited states. The comparison between our data and the calculations indicates a strong dominance of $l = 0$ transfer for the sum of the contribution of the two $3/2^+$ levels and the value obtained for the total spectroscopic factor is $S'_1 + S'_2 = 0.21$ [15]. Using an alternate set of optical parameters from a compilation[11] led to the same spectroscopic factor value showing the stability of the result with respect to variation of the optical parameters. Although no peak is seen in Figure 2 for the $1/2^-$ level ($E_x = 6.429$ MeV, $\Gamma = 280$ keV) due to its large total width, it is possible to derive an upper limit for the spectroscopic factor of $S'_3 \lesssim 0.15$ assuming an $l = 1$ transfer (see Figure 2). The important result of this analysis is that the contribution of these two $3/2^+$ resonances to the destruction rate of ^{18}F *cannot* be neglected. Sources of uncertainty on spectroscopic factors are statistics (negligible), normalization to elastic scattering data ($\approx 10\%$) but the major one is the DWBA method itself. This last one is difficult to evaluate but using statistics of ratios between the directly measured proton widths and the ones extracted from a DWBA analysis[12], we estimate this error to 25% (r.m.s.) for spectroscopic factors greater than 0.01. Larger deviation occurs for smaller spectroscopic factor due to the contribution of other mechanisms, but the high value of 0.21 provides a safe margin.




FIG. 2: Example of a reconstructed ^{19}F coincidence spectrum showing the two $3/2^+$ levels of astrophysical interest around 6.5 MeV of excitation energy. The bold line shows the calculated contribution of the $1/2^-$ level at 6429 keV for $S'_3 = 0.15$ (see text). This spectrum is limited at low energy by the coincidence condition (no ^{19}F breakup) and at high energy ($E_X \approx 7.3$ MeV) by the proton energy below electronic threshold.




FIG. 3: Comparison between the experimental angular distribution and DWBA calculations for different transferred angular momentum. The vertical error bars are only statistical whereas the horizontal ones are the angular width of each strip as seen from the target.

To show the improvement brought by this experiment, we present in Figure 4 the rates calculated in the same

way as in Ref. [3] except that we use new spectroscopic factors obtained from this experiment and the new value for the 330 keV resonance strength[5] ($\omega\gamma = 1.48 \pm 0.46$ eV). Assuming the equality of spectroscopic factors between analog levels one obtains $S_1 + S_2 = S'_1 + S'_2 = 0.21$ and $S_3 = S'_3 < 0.15$ where S'_1 , S'_2 and S'_3 correspond to the $E_x, J^\pi = 6.497$ MeV, $3/2^+$; 6.528 MeV, $3/2^+$; 6.429 MeV, $1/2^-$ levels in ^{19}F while S_1 , S_2 and S_3 correspond to the $E_x, J^\pi = 6.419$ MeV, $3/2^+$; 6.449 MeV, $3/2^+$; 6.437 MeV, $1/2^-$ analog levels in ^{19}Ne , respectively. Taking into account the moderate uncertainty on the energy calibration, the data favors a dominant contribution from the 6.528 MeV level i.e. $S'_1 \approx 0$ and $S'_2 \approx 0.2$. However the assignment of the two analog levels in ^{19}Ne is not firmly established. We cannot exclude inversion or mixing of these two $3/2^+$ levels. Hence to be conservative, for the calculation of the rate, we allow the individual spectroscopic factors to take any value compatible with the $S_1 + S_2 = 0.21$ and $S_3 < 0.15$ constraints. To show the improvement provided by this experiment, we calculate upper and lower rate limits by setting $S_1 = 0$, $S_2 = 0.21$ (no inversion), $S_3 = 0.15$ and $S_1 = 0.21$, $S_2 = 0$ (inversion), $S_3 = 0$ respectively. Here we neglect the error associated with the spectroscopic




FIG. 4: Present limits to the $^{18}\text{F}(p,\alpha)^{15}\text{O}$ reaction rate, calculated from this work and the new 330 keV resonance strength[5] (hatched area), compared to the previous situation[3] (CHJT00). All rates are normalized to the Coc et al. nominal rate[3] used in the more recent nova calculations. The Wiescher and Kettner rate[13] (WK82) used in the earliest nova calculations[1] is also displayed for comparison.

factor extraction discussed above. As a consequence, the rate uncertainty is reduced by a factor of ≈ 5 (Figure 4) in the temperature range of novae. This is mainly due to the reduced contribution of the $1/2^-$ level while the impact on the rate of the new contribution of the two $3/2^+$ levels is more important at low temperatures. It is worth noticing that these new rate limits still encompass the nominal rate[3] previously used in the most recent gamma-ray flux calculations. Since the repartition of spectroscopic factors between the two $3/2^+$ levels is not settled, the choice $S_1 = 0.1$ and $S_2 = 0.1$ for the nominal rate[3] is still acceptable. However, the new strength[5] of the resonance at 330 keV has to be used for the rate calculation.

This work has been supported by the European Community-Access to Research Infrastructure action of the Improving Human Potential Program, contract N HPRI-CT-1999-00110. One of us (P.L.) is a Research Director of the National Fund for Scientific Research, Brussels. We thank Alan Shotton and his team for allowing us to use the LEDA and LAMP detectors from the Louvain-La-Neuve and Edinburgh collaboration.

-
- [1] J. Gómez-Gomar, et al., *MNRAS* **296**, 913 (1998).
 - [2] M. Hernanz, et al., *Astrophys. J.* **526**, L97 (1999).
 - [3] A. Coc, et al., *Astron. Astrophys.* **357**, 561 (2000).
 - [4] J.-S. Graulich, et al., *Phys. Rev.* **C63**, 011302 (2001).
 - [5] D.W. Bardayan, et al., *Phys. Rev. Lett.* **89**, 262501 (2002).
 - [6] M. Cogneau, et al., *Nucl. Inst. and Meth.* **A420**, 489 (1999).
 - [7] M.E.O de López, et al., *Nucl. Phys.* **51**, 321 (1964).
 - [8] T. Davinson, et al., *Nucl. Inst. and Meth.* **A454**, 350 (2000).
 - [9] D. R. Tilley, et al., *Nucl. Phys.* **595**, 1 (1990).
 - [10] I. J. Thompson, *Computer Physics Report* **7**, 167, (1988).
 - [11] C. M. Perey, et al., *Atomic Data and Nuclear Data Tables* **17**, 1 (1976).
 - [12] J. Vernotte, et al., *Phys. Rev.* **C41**, 1956 (1990).
 - [13] Wiescher M. and Kettner, K.U., *Astrophys. J.* **263**, 891 (1982).
 - [14] N. de Séreville, et al., *Proceedings of the International Conference on Classical Nova Explosion*, Sitges, Spain, 20-24 May 2002, AIP, 2002., nucl-ex/0206014.
 - [15] Preliminary results from Kozub et al. quoted in a very recent paper[5] are in agreement with our value[14].